BULLETIN OF ENVIRONMENTAL SCIENCE AND TECHNOLOGY

VOL.1-NO.1-2026

P ISSN: xxxx-xxx

E ISSN: xxxx-xxx

ENVIRONMENTAL MICROBIOLOGY AND PUBLIC HEALTH

Hasan Ali

BS Student, Department of English, University of Okara

ABSTRACT

Environmental microbiology is an interdisciplinary field that examines microorganisms and their interactions with water, soil, air, wastewater, food, plants, animals, and human populations. Its importance to public health has increased substantially because environmental microbial communities can influence the emergence, persistence, and transmission of infectious diseases, antimicrobial resistance (AMR), and other biological hazards. Microorganisms present in environmental systems may be beneficial by supporting nutrient cycling, biodegradation, ecosystem stability, and biological remediation; however, pathogenic and opportunistic microorganisms can create significant risks when environmental conditions facilitate their survival and dissemination. Contaminated drinking water, untreated wastewater, poorly managed solid waste, agricultural runoff, healthcare waste, and polluted recreational waters can provide pathways through which bacteria, viruses, fungi, parasites, and antimicrobial-resistance determinants reach human populations. Recent research increasingly emphasizes the environmental dimension of AMR because wastewater, soil, sediments, and other ecological compartments can function as reservoirs of antimicrobial-resistant bacteria and resistance genes. The World Health Organization has specifically highlighted wastewater, environmental surveillance, sanitation, and pollution control as important components of AMR prevention and public-health protection. Environmental microbiology also contributes to public-health surveillance through microbial monitoring and wastewater-based environmental surveillance. Conventional microbiological culture remains valuable for identifying viable organisms and determining antimicrobial susceptibility, while molecular methods such as polymerase chain reaction (PCR), quantitative PCR (qPCR), sequencing, and metagenomics provide increasingly detailed information about microbial communities and resistance genes. These approaches can support early detection of pathogens, identification of contamination sources, monitoring of environmental AMR, and assessment of wastewater-treatment performance. WHO guidance published in recent years recognizes wastewater and environmental surveillance as an important component of pathogen monitoring and provides specific guidance concerning AMR surveillance.

KEYWORDS: *Environmental microbiology, public health, microorganisms, pathogenic bacteria, wastewater, water quality, antimicrobial resistance, environmental surveillance, One Health, microbial contamination, sanitation, infectious diseases*.

INTRODUCTION

Environmental microbiology has become an increasingly important area of scientific research because microorganisms are fundamental components of virtually every ecosystem and are closely connected with human health. Bacteria, viruses, fungi, protozoa, archaea, and other microorganisms inhabit soil, water, air, sediments, plants, animals, wastewater, food systems, and human-associated environments. Many of these microorganisms are harmless or beneficial, contributing to nutrient cycling, decomposition, soil fertility, wastewater treatment, food production, and ecosystem functioning. At the same time, certain microorganisms can cause infectious diseases or act as reservoirs of genes and biological characteristics that create risks to human health. Consequently, understanding microorganisms in environmental settings is essential not only for ecological science but also

for disease prevention, environmental management, and public-health protection.The relationship between the environment and human health is particularly evident in water and sanitation. Microbial contamination of drinking water can expose populations to enteric pathogens, while inadequate wastewater treatment can allow microorganisms and microbial genetic material to enter rivers, lakes, groundwater, agricultural environments, and coastal waters. Environmental contamination can therefore create multiple pathways of human exposure, including drinking contaminated water, consuming contaminated food, contact with recreational water, occupational exposure, and indirect exposure through agricultural products. Environmental microbiology provides the scientific tools required to detect and characterize these hazards and to understand their persistence and movement through ecological systems.

Wastewater is particularly important because it receives microorganisms and chemical substances originating from households, healthcare facilities, agriculture, industries, and other human activities. Wastewater may contain pathogenic microorganisms, antimicrobial-resistant bacteria, antibiotic residues, resistance genes, nutrients, heavy metals, and other pollutants. Treatment processes can substantially reduce microbial contamination, but treatment efficiency varies according to technology, operating conditions, pollutant characteristics, and local infrastructure. Inadequate treatment or uncontrolled discharge can consequently increase environmental exposure. WHO has emphasized the importance of wastewater management and environmental action in addressing AMR and protecting public health One of the most significant developments in contemporary environmental microbiology is the increasing recognition of antimicrobial resistance as an environmental and public-health problem. AMR occurs when microorganisms develop or acquire characteristics that allow them to survive exposure to antimicrobial medicines. Although antimicrobial resistance has traditionally been considered primarily a clinical problem, evidence increasingly demonstrates that environmental systems participate in the maintenance and dissemination of resistant microorganisms and resistance genes. Wastewater from hospitals and communities, pharmaceutical manufacturing effluents, animal production systems, agricultural runoff, sewage, and contaminated soils can all contribute to environmental exposure to antimicrobial residues and resistant microorganisms. WHO has noted that environmental pollution can influence the development, transmission, and spread of AMR. (World Health Organization)Environmental microbiology is important in this context because microbial communities do not exist as isolated organisms.

They form complex ecosystems in which microorganisms interact with one another and with physical and chemical environmental conditions. Genetic material can move between microorganisms through horizontal gene transfer, potentially allowing resistance genes to spread among bacterial populations. Environmental factors such as antibiotic residues, heavy metals, disinfectants, nutrient availability, temperature, pH, and microbial community composition may influence microbial survival and gene transfer. Recent reviews have emphasized the role of environmental microbiomes as reservoirs and transmission pathways

for antimicrobial-resistance genes and have identified environmental surveillance as an important component of One Health The One Health framework provides an important conceptual basis for understanding environmental microbiology and public health. One Health recognizes that human health, animal health, plant health, and environmental health are interconnected. Microorganisms and resistance determinants can move among these compartments through water, food, soil, animals, waste, and human activities. For example, resistant bacteria may originate in healthcare or agricultural environments, enter wastewater or soil, persist in environmental reservoirs, and subsequently reach humans through water, food, direct contact, or other pathways. Recent literature therefore emphasizes coordinated surveillance and intervention across human, animal, and environmental sectors rather than treating each sector separately.Environmental microbiology also has an important role in infectious-disease surveillance. Microbial monitoring can identify contamination before or during disease outbreaks and can help public-health authorities understand the distribution of pathogens in communities. Wastewater-based surveillance has received considerable attention because wastewater can provide population-level information about microbial circulation. WHO's current environmental-surveillance guidance includes pathogen-specific and AMR-focused approaches, demonstrating the increasing institutional importance of environmental microbiology in public-health surveillanc

Technological developments have significantly expanded the capacity of environmental microbiology. Traditional culture-based techniques allow researchers to isolate viable microorganisms and determine their phenotypic characteristics. Microscopy, biochemical testing, selective culture, and antimicrobial-susceptibility testing remain important tools. Molecular techniques have added new capabilities by allowing researchers to detect microorganisms and genes that may be difficult or impossible to culture using conventional laboratory methods. PCR and qPCR can detect specific organisms or genetic markers, whereas sequencing and metagenomics can characterize microbial communities and identify multiple resistance determinants simultaneously. Metagenomic approaches are particularly valuable because environmental samples contain complex microbial communities. Instead of examining only one organism at a time, researchers can analyze genetic material from entire microbial communities. This approach can reveal the composition of environmental microbiomes, identify resistance genes, detect potential pathogens, and investigate relationships between microorganisms and environmental conditions. Nevertheless, molecular detection does not always demonstrate that an organism is viable or capable of causing disease. Therefore, environmental microbiological research benefits from integrating culture-based, molecular, ecological, and epidemiological evidence.Water quality represents another major connection between environmental microbiology and public health. Safe water requires control of microbial pathogens as well as chemical contaminants. Indicator organisms, including coliform bacteria and Escherichia coli, are commonly used to assess fecal contamination and evaluate water-quality conditions.

However, microbial indicators cannot represent every possible pathogen, and modern surveillance increasingly incorporates molecular approaches and pathogen-specific testing. Poor sanitation and contaminated water remain important public-health concerns, particularly where wastewater treatment, drinking-water infrastructure, waste management, and hygiene services are inadequate.Soil is also an important environmental microbial reservoir. Soil contains extremely diverse microbial communities that participate in nutrient cycling and organic-matter decomposition. However, soil can become contaminated with pathogens, antimicrobial-resistant bacteria, pharmaceutical residues, heavy metals, pesticides, sewage, and agricultural waste. Agricultural practices may create connections between soil microbiomes, livestock, crops, water systems, and human populations. Manure application, irrigation with contaminated water, and inadequate waste management can potentially introduce microorganisms and resistance determinants into agricultural environments. Air and airborne particles also contribute to environmental microbial exposure. Microorganisms can become associated with aerosols, dust, droplets, bioaerosols, and particulate matter.

RESEARCH OBJECTIVES

1. To examine the role of environmental microorganisms in public health, with particular emphasis on microbial contamination of water, wastewater, soil, food, air, and other environmental compartments.
2. To analyze the relationship between environmental microbial contamination and antimicrobial resistance, including the role of wastewater, agricultural activities, healthcare facilities, and environmental reservoirs in the dissemination of resistant microorganisms and resistance genes.
3. To evaluate the importance of environmental microbiological surveillance and preventive strategies for reducing infectious-disease risks and supporting sustainable public-health protection under the One Health framework.

RESEARCH QUESTIONS

1. What role do environmental microorganisms and microbial contamination play in public health?
2. How do environmental conditions and human activities contribute to the emergence and dissemination of antimicrobial-resistant microorganisms?
3. How can environmental microbiological surveillance, wastewater management, sanitation, and One Health interventions contribute to improved public-health protection?

LITERATURE REVIEW

Environmental microbiology has increasingly become central to understanding the connections between environmental quality and human health. Contemporary research demonstrates that microorganisms are not merely isolated causes of disease but components

of complex environmental systems in which microbial communities interact with humans, animals, plants, pollutants, and ecological conditions. The microbiome perspective has particularly expanded understanding of these interactions. A 2023 review of the One Health microbiome relationship reported that microbial communities contribute to interactions among humans, animals, and environmental systems and that resistance genes within microbial communities can contribute to antimicrobial resistance across interconnected ecosystems. These findings support an integrated approach in which environmental microbial processes are considered alongside clinical and epidemiological information. A major theme in recent literature is the environmental dimension of antimicrobial resistance. Martak et al. (2024) describe environmental reservoirs as important sources of resistant bacteria, resistance genes, and selective agents, emphasizing the contribution of sewage, manure, industrial waste, antibiotics, biocides, and heavy metals to environmental AMR Similarly, Yuan et al. (2024) reviewed transmission pathways connecting environmental AMR with clinical settings and highlighted horizontal gene transfer and environmental pressures as mechanisms that may facilitate the movement of resistance from environmental microorganisms toward clinically relevant bacteria. These findings indicate that AMR surveillance needs to extend beyond hospitals and clinical samples to include environmental reservoirs.

Environmental surveillance of wastewater can therefore provide information about microbial hazards at a population level. WHO's environmental-surveillance framework identifies wastewater as a useful matrix for monitoring pathogens and antimicrobial resistance when appropriate sampling, laboratory, interpretation, and quality-control procedures anWastewater surveillance should nevertheless be interpreted carefully because detection of genetic material does not necessarily mean that viable pathogens are present or that transmission will occur. Environmental microbiology has also become important in understanding the ecological processes behind AMR. The environmental resistome refers broadly to the collection of antimicrobial-resistance genes present within microbial communities. Recent research emphasizes that soil, water, sediments, wastewater, and other environmental compartments can contain diverse resistance determinants. Environmental microbial communities can potentially facilitate gene exchange through mobile genetic elements and horizontal gene transfer. A recent review emphasizes that environmental microbiomes can act as reservoirs and transmission vectors of resistance genes and argues that environmental surveillance is particularly important in regions where monitoring infrastructure remains limited. (PubMed)Another important research direction concerns the relationship between microbial diversity and resistance dissemination.

This perspective demonstrates that environmental microbiology cannot be reduced to counting pathogenic organisms. Researchers increasingly need to understand microbial community structure and ecological interactions because these factors may influence the persistence and transmission of resistance.Agriculture is another important component of the environmental microbiology-public health relationship. Animal manure, agricultural runoff,

irrigation water, and soil amendments can introduce microorganisms and antimicrobial residues into environmental systems. Resistant organisms may subsequently move between animals, soil, water, crops, wildlife, and humans. Research using the One Health perspective has documented interconnected reservoirs in humans, animals, food, and the environment. Such findings reinforce the need for surveillance systems that integrate agricultural, environmental, veterinary, and public-health information.Healthcare facilities are similarly important because they can release pathogens, resistant bacteria, antimicrobial residues, disinfectants, and other biological materials into wastewater systems. Infection prevention and control therefore cannot be separated completely from environmental management. WHO's 2024 global report identifies water, sanitation, hygiene, waste management, infection prevention and control, and AMR as interconnected areas of health-system protecti Environmental microbiological monitoring can complement clinical surveillance by identifying microbial hazards in wastewater and healthcare environments.

Improved sanitation reduces human exposure to fecal microorganisms, while safe water and hygiene practices reduce transmission opportunities. One Health research has increasingly connected WASH improvements with reductions in infection risk and reduced demand for antimicrobial treatment. (PubMed) Environmental microbiology provides the laboratory evidence necessary to evaluate the effectiveness of these interventions. Recent research also emphasizes the need for standardized environmental AMR risk assessment. The complexity of environmental ecosystems makes it difficult to translate the detection of a resistance gene into a specific human-health risk. A 2024 review on environmental AMR indicators highlighted the need to integrate ecological data, microbiome information, risk assessment, and emerging technologies to create more useful environmental AMR early-warning systemsTherefore, future environmental microbiology should combine microbiological detection with epidemiological, ecological, and risk-assessment approaches.Taken together, the literature demonstrates that environmental microbiology has moved from a primarily ecological discipline toward a major component of public-health science. Water, wastewater, soil, food, animals, healthcare systems, industry, and human communities are interconnected microbial environments. The strongest contemporary research supports One Health surveillance, integrated microbial monitoring, improved sanitation, wastewater treatment, responsible antimicrobial use, and stronger environmental policies.

METHODOLOGY

This study adopts a mixed-methods-oriented environmental public-health approach, combining microbiological, environmental, and public-health perspectives. The research framework is designed to examine microbial hazards in environmental systems and their potential implications for human health. The approach incorporates evidence from environmental microbiology, epidemiology, wastewater science, antimicrobial-resistance research, and One Health literature. Environmental matrices considered within the framework include drinking water, wastewater, surface water, soil, agricultural environments, healthcare

waste, and other settings where microbial contamination can occur. The study emphasizes systematic interpretation of environmental microbial indicators rather than considering individual microorganisms independently. This approach is consistent with current international recognition that environmental factors contribute to AMR development and transmission.The analytical component focuses on identifying relationships among environmental contamination, microbial hazards, antimicrobial resistance, sanitation conditions, and public-health risks. Where microbiological data are available, variables may include microbial counts, pathogen detection, resistance profiles, resistance genes, environmental conditions, wastewater-treatment characteristics, and contamination sources. Qualitative evidence from recent scientific literature and international guidance is used to contextualize quantitative microbiological observations. The mixed perspective is particularly appropriate because environmental microbiology involves both measurable laboratory parameters and complex ecological processes that cannot always be explained through numerical measurements alone.

RESEARCH DESIGN

The study uses a descriptive and analytical research design. The descriptive component identifies the major categories of environmental microorganisms, contamination sources, environmental reservoirs, exposure pathways, and public-health implications. The analytical component examines relationships between environmental factors and microbial risks, particularly the relationship between wastewater contamination, antimicrobial residues, resistant microorganisms, and resistance genes. The design allows environmental microbiological information to be organized systematically while also examining factors that may contribute to microbial persistence and dissemination.A structured literature-based research component is incorporated into the design to synthesize recent evidence from peer-reviewed studies and authoritative public-health organizations. Priority is given to recent evidence from 2023–2026, supplemented where necessary by established scientific literature and international guidance. Key themes include environmental pathogens, wastewater microbiology, AMR, environmental surveillance, One Health, sanitation, microbial risk assessment, and microbiological monitoring. WHO's recent environmental-surveillance and AMR documents provide an important framework for interpreting environmental microbiology within

DATA ANALYSIS

Data analysis in environmental microbiology should begin with systematic organization and quality assessment of microbiological observations. If environmental samples are collected, samples can be categorized according to source, including drinking water, wastewater, surface water, soil, agricultural runoff, healthcare wastewater, and other relevant environmental compartments. Microbial measurements can then be summarized using descriptive statistics such as frequencies, percentages, means, medians, standard deviations, minimum values, and maximum values. Where appropriate, microbial concentrations can be

expressed using standardized units such as colony-forming units per volume or mass of sample. Molecular results may be presented according to detection frequency, gene abundance, relative abundance, or quantitative gene copies. These measurements can then be compared across environmental locations, sampling periods, and contamination sources. For AMR analysis, resistance profiles can be summarized by antimicrobial class and microorganism, while the presence of resistance genes can be compared among different environmental matrices. The second stage of analysis examines relationships among environmental and microbiological variables. Correlation or regression techniques may be used where the data meet relevant statistical assumptions. For example, researchers can investigate whether microbial contamination is associated with wastewater-treatment characteristics, environmental conditions, population density, antibiotic residues, or other potential explanatory variables.

FINDINGS

The findings indicate that environmental microbiology represents a fundamental component of modern public-health protection because environmental microorganisms can simultaneously provide ecological benefits and create biological hazards. Environmental systems contain highly diverse microbial communities that participate in nutrient cycling, decomposition, pollutant transformation, and ecosystem stability. However, environmental contamination can introduce pathogenic and opportunistic microorganisms into water, soil, food, wastewater, and other exposure pathways. The evidence reviewed in this study indicates that wastewater is particularly important because it can receive microorganisms and genetic material from households, hospitals, agriculture, industry, and other human activities. Environmental surveillance can therefore provide valuable information about microbial hazards circulating within communities. Current WHO guidance specifically recognizes wastewater and environmental surveillance as tools that can support pathogen and AMR monitoring. (World Health Organization) A major finding concerns the relationship between environmental contamination and antimicrobial resistance. The reviewed evidence indicates that environmental systems can function as reservoirs of resistant microorganisms and resistance genes. Wastewater, sewage sludge, agricultural environments, surface water, sediments, and contaminated soils can contain antimicrobial-resistant organisms and resistance determinants.

CONCLUSION

Environmental microbiology has become an essential component of public-health science because environmental microbial communities influence infectious-disease risks, antimicrobial resistance, water quality, sanitation, wastewater safety, and ecosystem health. The evidence reviewed in this study demonstrates that environmental systems can serve simultaneously as beneficial microbial ecosystems and reservoirs of potentially harmful

microorganisms and resistance determinants. Wastewater, soil, surface water, agricultural environments, healthcare facilities, and industrial systems are particularly important interfaces between microorganisms and human populations. Environmental surveillance can strengthen public-health monitoring by providing information that complements clinical and laboratory surveillance.The study further concludes that antimicrobial resistance must be addressed through an integrated environmental and One Health framework. Resistant microorganisms and resistance genes can move among humans, animals, food, water, soil, and other environmental compartments. Consequently, effective public-health protection requires improvements in wastewater treatment, sanitation, hygiene, healthcare waste management, pharmaceutical pollution control, agricultural practices, environmental surveillance, and responsible antimicrobial use. Recent WHO guidance and scientific literature strongly support greater attention to the environmental dimension of AMR. (World Health Organization) Strengthening environmental microbiology laboratories, surveillance systems, research capacity, and interdisciplinary collaboration can contribute to earlier identification of microbial hazards and more effective prevention strategies.

REFERENCES

Pepper, I. L., Gerba, C. P., & Gentry, T. J. (2015). Introduction to environmental microbiology. In Environmental microbiology (pp. 3-8). Academic Press.

World Health Organization. (2025). European health report 2024: keeping health high on the agenda: highlights.

. World Health Organization. (2025). Civil society engagement in the development of World Health Assembly resolutions and decisions: lessons and examples from the Seventy-seventh session. World Health Organization.

World Health Organization. (2024). COVID-19 epidemiological update, edition 174, 24 December 2024.

World Health Organization. (2024). COVID-19 epidemiological update, edition 174, 24 December 2024.

World Health Organization. (2025). State of the world's nursing 2025: investing in education, jobs, leadership and service delivery. World Health Organization.

Gu, X., Watson, C., Agrawal, U., Whitaker, H., Elson, W. H., Anand, S., ... & de Lusignan, S. (2024). Postpandemic sentinel surveillance of respiratory diseases in the context of the world health organization mosaic framework: Protocol for a development and evaluation study involving the english primary care network 2023-2024. JMIR public health and surveillance, 10(1), e52047.